

# NONLINEAR PHENOMENA OF FRÖHLICH PHONONS IN BIOLOGICAL MEDIA

Anatol Rotaru, Nelu Ciobanu, and V. Z. Tronciu

*State University of Medicine and Pharmaceutics „Nicolae Testemitanu,  
Technical University of Moldova  
tronciu@mail.utm.md*

(Received October 21, 2013)

## Abstract

Nonlinear cooperative stationary phenomena are studied for the interaction of Bose-condensed phonons with millimeter electromagnetic radiation in the biological media. The real and imaginary parts of the dielectric susceptibility and permeability, refraction and reflection indexes determined by Bose-condensed phonons are obtained. The possibility of occurrence of the polariton effect in biological objects is predicted.

## 1. Introduction

The investigation of the nonlinear cooperative phenomena in complex systems is one of the most important and up-to-date problems of the modern science. In recent years, interest in the investigation of the interaction of the millimeter range coherent electromagnetic radiation with medical-biological objects has attracted much attention due to many applications of millimeter waves in the applied medicine, biotechnology, and agriculture [1–8].

The concept of generation of phonons in alive media was proposed by Fröhlich [9]. He theoretically suggested that biological systems can generate collective vibrational modes (phonons) in the GHz frequency range and these phonons might play a basic role in the active biological systems. The idea is based on the assumption that, if the generated energy is supplied to the phonon at a rate greater than a certain critical value, then the phenomenon of Bose condensation to the lowest excitation of a single mode can occur. This kind of condensation that occurs at the lowest excitation mode can serve as a means for energy storage as well as a channel for specific bio-processes, such as cell division or macromolecule synthesis. This idea stimulated further investigations in the domain with an aim of studying and predicting new biophysical phenomena and their effective use in biomedicine [10–15].

Although the number of experimental and theoretical papers in the field continuously grows, up to the present there is no clear understanding of the mechanism of generation and interaction of the millimeter radiation with the biological media. Thus, a consistent theory of the low-intensity millimeter wave interaction with biological media has not been elaborated, and a lot of cooperative nonlinear phenomena have not yet been studied. It is well known that, due to the metabolism processes in the organisms, the excitation of the polarizing coherent waves is possible in a frequency range of  $10^{11}$ - $10^{12}$  Hz. These oscillations are supposed to cover parts of biological membranes and, as was demonstrated, it can be equivalent to Bose-Einstein condensation of phonons in these media. For example, Davydov investigated the possibility of

the collective state excitation in one-dimensional molecular structures and in double-helix protein molecules in the form of solitons [16, 17], which was further developed for dissipative structures [18]. Nonlinear, collective, and stochastic phenomena of Bose-type quasiparticles in condensed media and, in particular, the nonlinear dynamics of the immune system interaction with the bilocal cancer tumor were developed in [19]. A new simple model of local interaction between two spaced cancer cells and cytotoxical T-lymphocytes is proposed.

This paper is focused on the investigation of nonlinear cooperative phenomena at the interaction of a Bose-condensed phonon with the millimeter electromagnetic radiation generated in a biological object. We start in Section II with a Hamiltonian of the interaction of millimeter electromagnetic radiation with coherent phonons in the square form for the low concentrations of Bose-condensed phonons. The collective nonlinear properties of the coherent phonons are determined, in particular, by the phonon-phonon interaction pattern, the energetic spectrum of the coherent phonons and their interaction with the non-condensed quasiparticles. Section III concerns the study of the real and imaginary parts of dielectric susceptibility and permeability, refraction and reflection indexes determined by Bose-condensed phonons. This study represents a major interest in modern biophysics. In Section IV, the numerical results on calculations of system characteristics are presented. The possibility of occurrence of a polariton is also discussed. Conclusions are given in section V.

## 2. Model

In what follows, based on the Fröhlich idea that coherent phonons excited in biological objects turn into coherent internal photons and form a self-correlated interval millimeter electromagnetic field, we derive a set of equations that describe the evolution of these photons and phonons. As was previously demonstrated, the coherent state of the phonon in biological objects can be similar to the coherent state of the exciton in condensed media [20–23]. These coherent states of the elementary excitations can appear in different regimes; i.e., in the regime of ultrashort pulses existing for a time period shorter than the relaxation times as well as in conditions of Bose-Einstein condensation for a time period longer than the relaxation times, but shorter than the quasiparticle life time. In both cases, the corresponding elementary excitations are characterized by a certain phase, wave vector, and macroscopic value of the single-particle state; that is, the number of excitations in condensate  $N_V \sim V$ , where  $V$  is the system volume.

The Hamiltonian of the system has the form

$$H = \sum_{\vec{k}} \hbar\Omega a_{\vec{k}}^+ a_{\vec{k}} + \frac{1}{8\pi} \sum_{\vec{k}} \int (\varepsilon E_{\vec{k}}^2 + \mu H_{\vec{k}}^2) dv_{\vec{k}} + \frac{1}{2V} \sum_{\vec{k}_1, \vec{k}_2, \vec{k}'_1, \vec{k}'_2} g(\vec{k}_1 - \vec{k}'_1) \delta_{kp}(\vec{k}_1 + \vec{k}_2, \vec{k}'_1 + \vec{k}'_2) a_{\vec{k}_1}^+ a_{\vec{k}_2}^+ a_{\vec{k}'_1} a_{\vec{k}'_2} - \sum_{\vec{k}} d_{\vec{k}} (E_{\vec{k}}^+ a_{\vec{k}}^+ + E_{\vec{k}}^- a_{\vec{k}}^-), \quad (1)$$

where  $\hbar\Omega$  is the dipole-active phonon energy;  $E_{\vec{k}}^- = E_{\vec{k}}^+ + E_{\vec{k}}^-$ ,  $H_{\vec{k}}^- = H_{\vec{k}}^+ + H_{\vec{k}}^-$  are the intensities of the electrical and magnetic fields, respectively;  $E_{\vec{k}}^{\pm}$  and  $H_{\vec{k}}^{\pm}$  are the positive or negative frequency parts of the variable electromagnetic field;  $a_{\vec{k}}^+$  ( $a_{\vec{k}}^-$ ) are the operators of creation (annihilation) of the dipole-active phonons that satisfy the commutation relation  $[a_{\vec{k}}, a_{\vec{k}'}^+] = \delta_{kk'}$ ,  $[a_{\vec{k}}, a_{\vec{k}}^+] = 0$ ;  $\varepsilon$  and  $\mu$  are the dielectric and magnetic permeability of the biological medium;  $g(\vec{k})$  is the phonon-phonon interaction constant; and  $d_{\vec{k}}$  is the dipole moment of the transition into phonon state.

In Hamiltonian (1), both the field and the Bose-condensed phonons are supposed to have the same wave vector  $\vec{k} \neq 0$ , polarization, and phase. The amplitudes are considered to be macroscopically large, i.e., proportional to system effective volume  $V$ . Note that the effects of spatial dispersion were neglected being insignificant in the actual spectrum range. Thus, we consider the case of phonon mass  $m \approx \infty$ .

The Heisenberg motion equation for the operator  $a_{\vec{k}}$  has the form

$$i \frac{da}{dt} = \Omega a + \frac{g}{\hbar} \frac{|a|^2}{V} a + \frac{d}{\hbar} E^+ - i\gamma a, \quad (2)$$

where the wave vector indexes are omitted. Note that we phenomenologically introduced the term  $i\gamma a$ , which takes into account the Bose-condensed phonons leaving from condensed state, in (2). On the other hand, the attenuation term can be considered strictly within the framework of the quantum theory of attenuations and fluctuations using the Fokker-Plank master equation [24].

The polarization of the biological media is given by

$$P^+ = -\frac{1}{V_o} \frac{\partial H}{\partial E^-}, \quad P^- = -\frac{1}{V_o} \frac{\partial H}{\partial E^+} \quad (3)$$

where  $V_o = \frac{V}{N_d}$ , and  $N_d$  is the atomic dipole number. Taking account Hamiltonian (1) and (3),

we obtain the following form for polarization

$$P^+ = \frac{da}{V_o}, \quad P^- = \frac{da^+}{V_o}. \quad (4)$$

We assume that the Fröhlich electromagnetic millimeter field is spread along the  $x$  axis. Thus, the equation of the positive frequency component of the electromagnetic field is equivalent to

$$\frac{\partial^2 E^+}{\partial x^2} - \frac{1}{c^2} \frac{\partial^2 E^+}{\partial t^2} = \frac{4\pi}{c^2} \frac{\partial^2 P^+}{\partial t^2} \quad (5)$$

where  $c$  is the light velocity in the biological media.

Equations (2) and (5) completely describe the spatial-temporal evolution of the Bose-condensed phonons in the biological media, taking into account self-interaction and the interaction with the Fröhlich field.

We write the macroscopic amplitudes of the phonons and the field in the form of modulated plane waves with frequency  $\omega$  and wave vector  $k$

$$\begin{aligned} a(x,t) &= \sqrt{V} \tilde{a}(x,t) e^{-i\omega t + ikx}, \\ E^+(x,t) &= \sqrt{V} \tilde{E}(x,t) e^{-i\omega t + ikx}, \end{aligned} \quad (6)$$

where  $\tilde{a}(x,t)$  and  $\tilde{E}(x,t)$  are the slowly-varying amplitudes satisfying the following conditions

$$\begin{aligned} \left| \frac{\partial \tilde{a}}{\partial t} \right| &\ll \omega |\tilde{a}|, & \left| \frac{\partial \tilde{a}}{\partial x} \right| &\ll k |\tilde{a}|, \\ \left| \frac{\partial \tilde{E}}{\partial t} \right| &\ll \omega |\tilde{E}|, & \left| \frac{\partial \tilde{E}}{\partial x} \right| &\ll k |\tilde{E}|. \end{aligned}$$

Taking into account the above assumption, equation of motion (2) is transformed in

$$\frac{d\tilde{a}}{dt} = i \left[ \Delta + i\gamma - \frac{g}{\hbar} n \right] \tilde{a} + i \frac{d}{\hbar} \tilde{E}, \quad (7)$$

where  $\Delta = \omega - \Omega$  is the resonance detuning between the Fröhlich field frequency and the phonon frequency.  $n = |\tilde{a}|^2$  represents the concentration of the Bose-condensed phonons.

### 3. Stationary States

In this Section, we consider the steady states ( $d\tilde{a}/dt = 0$ ) of equation of motion (7). We obtain the dependence of amplitudes of phonons on photons

$$\tilde{a} = -\frac{d/\hbar}{\Delta - gn/\hbar + i\gamma} \tilde{E} \quad (8)$$

The dielectric susceptibility of the biological media is given by the relation  $\chi = P^+ / E^+$ . Taking into account (4) and (8) for the real and imaginary parts of the dielectric susceptibility, we obtain

$$\begin{aligned} \chi' &= -\frac{d^2}{V_0 \hbar} \frac{\Delta - gn/\hbar}{(\Delta - gn/\hbar)^2 + \gamma^2}, \\ \chi'' &= \frac{d^2 \gamma}{V_0 \hbar} \frac{1}{(\Delta - gn/\hbar)^2 + \gamma^2}. \end{aligned} \quad (9)$$

Thus, the dielectric permeability, that takes into account the phonon level contribution, is described by the expression

$$\varepsilon = \varepsilon_\infty + 4\pi\chi = \varepsilon' + i\varepsilon'', \quad (10)$$

where  $\varepsilon', \varepsilon''$  are the real and imaginary parts of the dielectric permeability

$$\begin{aligned} \varepsilon' &= \varepsilon_\infty - \frac{4\rho d^2}{V_0 \hbar} \frac{D - gn/\hbar}{(D - gn/\hbar)^2 + g^2} \\ \varepsilon'' &= \frac{4\rho d^2 g}{V_0 \hbar} \frac{1}{(D - gn/\hbar)^2 + g^2}. \end{aligned}$$

Here,  $\varepsilon_\infty$  is the phonon dielectric permeability that takes into account the dielectric permeability of all excitations besides phonons of the biological media. The real and imaginary parts of the permeability determine the frequency dispersion. From (10) we conclude that the dielectric permeability of the biological media depends significantly on Bose-condensed phonon density  $n$ . In what follows it is convenient to switch to the following dimensionless quantities

$$\begin{aligned} N &= \frac{g}{\gamma \hbar} n, \quad f = \frac{\tilde{E}}{E_s}, \quad \frac{1}{E_s^2} = \left( \frac{d}{\hbar} \right)^2 \frac{g}{\hbar \gamma^3}, \\ \Omega_0 &= \frac{4\pi d^2}{\hbar V_0}, \quad \delta = \frac{\Delta}{\gamma}. \end{aligned} \quad (11)$$

Taking account (11) from (10), we obtain

$$\begin{aligned} \varepsilon' &= \varepsilon_\infty - \frac{(\Omega_0 / \gamma)(\delta - N)}{[(\delta - N^2) + 1]}, \\ \varepsilon'' &= \frac{\Omega_0 / \gamma}{[(\delta - N)^2 + 1]}, \end{aligned}$$

$$N[(\delta - N)^2 + 1] = f^2. \quad (12)$$

For a plane wave, where the surfaces of the field components are planes, the dielectric permeability perpendicular to the spreading direction is related to refraction index  $\bar{n}$  and absorption index  $\kappa$  by the expression [25]

$$\varepsilon = \bar{n}^2 - \kappa^2 + 2i\bar{n}\kappa. \quad (13)$$

Thus, we can write the refraction and absorption indexes

$$\begin{aligned} \bar{n} &= \sqrt{\frac{\varepsilon' + \sqrt{\varepsilon'^2 + \varepsilon''^2}}{2}}, \\ \kappa &= \sqrt{\frac{-\varepsilon' + \sqrt{\varepsilon'^2 + \varepsilon''^2}}{2}} \end{aligned} \quad (14)$$

It is well known that reflection coefficient  $R$  is defined as the ratio of the time average energy flow reflected from the surface to the incident one. For the case of a perpendicular flow, the reflection coefficient can be written [25, 26]:

$$R = \frac{(\bar{n} - 1)^2 + \kappa^2}{(\bar{n} + 1)^2 + \kappa^2} \quad (15)$$

In what follows, consider that the electromagnetic field is uniformly distributed in the biological media ( $\partial\tilde{E}/\partial x = 0$ ). From (5) we obtain the equation for the field temporal evolution in the slowly-varying amplitude approximation:

$$\frac{d\tilde{E}}{dt} = i \left[ \frac{\omega^2 - c^2 k^2}{2\omega} - \frac{4\pi d^2}{V_o \hbar} \right] \tilde{E} + \frac{2\pi d \omega i}{V_o} \left[ 1 - \frac{2}{\omega} \left( \omega - \Omega - \frac{g}{\hbar} n \right) - \frac{2i\gamma}{\omega} \right] \tilde{a}. \quad (16)$$

In the stationary case  $\partial\tilde{E}/\partial t = 0$  and the dispersion limit, where the attenuation of the dipole-active Bose-condensed phonons can be neglected, from (16) we obtain

$$k^2 = \frac{\omega^2}{c^2} \left[ 1 + \frac{\Omega_o}{\Omega + \frac{g}{\hbar} n - \omega} \right]. \quad (17)$$

Note that equation (17) determines the dispersion law of the nonlinear polaritons; that is, the energy values of the elementary excitations in biological media at the Fröhlich radiation interaction with dipole-active Bose-condensed phonons can occur at the same value of the wave vector. The position and form of the polariton branches depend not only on the parameters of the phonons and the electrical field, but also on the stationary concentration of the Bose-condensed phonons. The concentration depends on the Fröhlich field intensity according to equation (12). Thus, the appearance of a polariton is caused by the intersection of the coherent photon dispersion curves with the Bose-condensed phonon for small wave vectors. Close to the intersection point, the energies and pulses of both excitations coincide and the bond between them becomes significant. As a result, neither photon nor phonon could be considered as independent elementary excitations in biological objects, but a new elementary excitation—polariton—appears in the biological objects.


#### 4. Numerical Results

As noted above, expression (17) coincides with that of dispersion law of the polariton in

the case of appearance of a soliton in the excitonic spectrum range [27]. Using the following dimensionless variables  $\bar{k} = k/k_o$ ,  $\bar{\omega} = \omega/(ck_o)$ ,  $\bar{\omega}_o = \Omega_o/(ck_o)$ , where  $ck_o = \Omega + gn/\hbar$ , from (17) we obtain the expression for dispersion law

$$\bar{k}^2 = \bar{\omega}^2 \left[ 1 + \frac{\bar{\omega}_o}{1 - \bar{\omega}} \right]. \quad (18)$$

According to (12), for small values of the field intensities, the dependence of the phonon concentration and the field is linear. While excitation increases, the phonon-phonon interaction processes play an important role. It is easy to observe that the relation between  $N$  and  $f$  is linear for  $\delta < \sqrt{3}$  (see Fig.1).




**Fig. 1.** Dependence of phonon concentration on field intensity for different values of resonance detuning: (1) for  $\delta = -5.0$ , (2) for  $\delta = 1.7$ , (3) for  $\delta = 5.0$ , (4) for  $\delta = 10.0$ , and (5) for  $\delta = 20.0$ .

For  $\delta > \sqrt{3}$ , the  $N(f)$  dependence is not linear any more and is characterized by a hysteresis region where three values of the dipole-active Bose-condensed phonon density correspond to one value of the Fröhlich field intensity. Figure 1 shows the dependence of phonon concentration on field intensity for different values of resonance detuning  $\delta$ . As it is shown in the figure, the growth of the field intensity leads to a jump-like increase in the Bose-condensed phonon density at  $\delta > \sqrt{3}$ . As the field decreases along the upper curve, a jump-like decrease in the phonon density occurs at  $\delta > \sqrt{3}$ . Thus, if the resonance detuning value is larger than the critical one  $\delta_c = \sqrt{3}$ , then the forward and backward alteration of the Fröhlich field intensity leads to jump-like changes in the phonon density and to the formation of an amplitude hysteresis loop in the  $N(f)$  dependence. The frequency hysteresis can be shown to occur also if one resonance detuning value corresponds to three values of the phonon density, values that are larger than the critical field intensity value  $f_c = (4/3)^{3/4}$ , one of which is unstable.


Note that the phenomena of nonlinear hysteresis present in the dependence of the Bose-condensed phonon density on Fröhlich field intensity have an effect on all dielectric functions: the real and imaginary parts of the dielectric susceptibility and permeability, the refraction, absorption, and reflection indexes. Figure 2 shows the dependences of the real (left) and imaginary (right) parts of the dielectric permeability of the biological object on field intensity

caused by Bose-condensed phonons for  $\varepsilon_\infty = 5.2, \bar{\omega} = 5.0$  and different values of the resonant detuning. If the resonance detuning values are smaller than the critical  $\delta_c$  value, then the real and imaginary parts of the dielectric permeability are single-valued functions of the field intensity. With an increase in the resonance detuning, the curves  $\varepsilon'(f)$ , and  $\varepsilon''(f)$  become deformed and hysteresis dependences  $\varepsilon'(f), \varepsilon''(f)$  appear at  $\delta > \delta_c$ .



**Fig. 2.** Dependences of the real and imaginary parts of the dielectric permeability on the amplitude of the field for  $\varepsilon_\infty = 5.2, \bar{\omega} = 5.0$  and the values of the resonant detuning similar to that in Fig. 1.

Figure 3 shows the dependences of the refraction, absorption, and reflection indexes on the field amplitude for different values of resonance detuning. From this figure, we conclude that for high values of the resonance detuning the  $n(f), \kappa(f)$  and  $R(f)$  dependences become more complicated. Thus, these functions exhibit a hysteresis behavior caused by the nonlinear dependence of the Bose-condensed phonon density on the coherent millimeter electromagnetic field amplitude if the resonance detuning exceeds the critical  $\delta_c$  value.



**Fig. 3.** Dependences of the refraction, absorption, and reflection indexes on field amplitude for the same parameters as in Fig. 2.

## 5. Conclusions

In this paper, we develop a model to describe the interaction of Bose-condensed phonons with millimeter electromagnetic radiation in the biological media. We obtain the real and imaginary parts of the dielectric susceptibility as well as the permeability, refraction, and reflection indexes determined by Bose-condensed phonons. The dielectric functions, as well as absorption and reflection coefficients, strictly depend on the coherent millimeter electromagnetic field amplitude and resonance detuning. For the values of the field intensity larger than the critical  $f_c$  value, a hysteresis appears in the system. The dispersion law shows that the interaction of Fröhlich radiation with dipole-active Bose-condensed phonons in biological media can lead to the occurrence of polaritons. We show that the position and form of the polariton branches depend on the stationary concentration of the Bose-condensed phonons. We mention that, for finite values of the phonon mass, equations (2) and (5) describe the electromagnetic field propagation and the generation of the coherent photons in biological media and can be represented by a system equivalent to the Ginzburg–Landau–Keldysh system. Neglecting the interaction of the coherent phonons with the thermostat  $g = 0$ , the above equations have soliton solutions. We believe that our work provides a good basis for future studies and, in particular, provides some pointers for more detailed investigations of the dynamics of Bose-condensed phonons with millimeter electromagnetic radiation in biological media.

**Acknowledgments.** V.Z.T. acknowledges the support from the Technical University of Moldova (projects 106 b/s, 13.820.05.08/GF) and from CIM - Returning Experts Programme.

## References

- [1] N. D. Deviatcov, and O. V. Betskii, *Biological Aspects of Low Intensity Millimeter Waves*, Moscow (1994).
- [2] O.V. Betskii and Y. G. Yaremenko, *Millimetr. Volny Biol. Med.* no. 1(11), 3 (1998).
- [3] A. Ramundo-Orlando, *J. Infrared Millimeter Terahertz Waves* 31, 1400 (2010).
- [4] X. -H. Li, J. -T. Tang, Y. -P. Liao, H. -K. Jin, J. -M. Zhou, G. -H. Wang, and H. Wang, *J. Clin. Rehab. Tissue Eng. Res.* 12, 663 (2008).
- [5] A. G. Pakhomov, Y. Akyel, O. N. Pakhomova, B. E. Stuck, and M. R. Murphy, *Bioelectromagnetics* 19, 393 (1998).
- [6] M. Markov, *Elec. Biol. And Med.* 26, 257 (2007).
- [7] W. -D. Li, W. Wang, J. -L. Chen, *World Chinese J. Digestology* 17, 3553-3557 (2009).
- [8] T. I. Usichenko, H. Edinger, V. V. Gizhko, C. Lehmann, M. Wendt, and F. Feyerherd, *Evidence-Based Complement. Altern. Med.* 3 (2), 201 (2006).
- [9] H. Fröhlich, *Phys Lett A* 26, 402 (1968); *Int. J. Quantum Chem.* 2, 641 (1968).
- [10] O. V. Betskii, D. V. Ghitsu, A. H. Rotaru, and N. V. Ciobanu, *The nonlinear dielectric permittivity during excitation of coherent phonons in biological objects. 13<sup>th</sup> Russian Symposium “Millimeter Waves in medicine and biology”, (Moscow, 2003), pp.118-121.*
- [11] M. Zhadobov, R. Saileau, V. Viè, M. Hindi, L. Le Coq, and D. Thouroude, *IEEE Trans. Microwave Theory Tech.* 54, 2534 (2006).
- [12] S. P. Sitko, *Phys. Alive.*1(1), 5 (1993).



- [13] V. P. Kalantaryan, Y. S. Babayan, E. S. Gevorgyan, S. N. Hakobayan, A. P. Antoyan, and P. O. Vardevayan, *Prog. Electromagn. Res. Lett.* 13, 1 (2010).
- [14] I. Szabo, J. Kappelmayer, S. I. Alekseev, and M. C. Ziskin, *Bioelectromagnetics* 27, 233 (2006).
- [15] I. A. Titushkin, V. S. Rao, W. F. Pickard, E. G. Moros, G. Shafirstein, and M.R. Cho, *Radiat. Res.* 172, 725 (2009).
- [16] A. S. Davydov, *J. Theor. Biol.* 66, 379 (1977).
- [17] A. S. Davydov, *Biology and Quantum Mechanics*. Pergamon, Oxford (1982).
- [18] H. Bolteraljer, and J. A. Tuszynski, *J. Biol. Phys.* 17, 41 (1989).
- [19] V. A. Zalozh, A. Kh. Rotaru, A. E. Barbaroshie, W. Van Driessche, and P. T. Frangopol, *J. Biol. Phys.* 21, 155 (1995).
- [20] A. H. Rotaru, and V. Tronciu, *JETP* 85, 971 (1997).
- [21] A. H. Rotaru, and C. Shura, *JETP* 86, 1132 (1998).
- [22] V. Tronciu, and A. H. Rotaru, *Phys. Solid State* 40, 1809 (1998).
- [23] A. H. Rotaru, and V. Tronciu, *Phys. Solid State B* 212, 383 (1999)
- [24] A. H. Rotaru, and G. D. Shibarshina, *Phys. Lett. A* 109, 292 (1985).
- [25] S. A. Moskalenko, A. Kh. Rotaru, and Yu. M. Shvera, *Teor. Mat. Fiz.* 75, 536 (1988).
- [26] L. D. Landau, and E. M. Lifshits, *The electrodynamics of continuous media*, Moscow (1982), (in Russian).
- [27] A. Kh. Rotaru, and V. A. Zalozh, *Optical Self-Organization of Excitons and Biexcitons in Semiconductors*, Shtiintsa, Kishinev (1990), (in Russian).